

## **Minimal-invasive Resektion eines Knochentumors im Bereich der temporoparietalen Schädelbasis**

### **Minimally invasive navigation-assisted removal of a bone tumor in the temporo-parietal skull base**

Jürgen Hoffmann<sup>1</sup>, Troitzsch, Dirk<sup>2</sup>, Dr.med. Westendorff, Carsten<sup>2</sup>, Priv.-Doz. Dr. Dr.  
Dammann, Florian<sup>3</sup>, Univ.-Prof. Dr. Dr. Reinert, Siegmars<sup>2</sup>

<sup>1</sup>Klinik und Poliklinik für Mund-, Kiefer- und Gesichtschirurgie, Universitätsklinikum  
Tübingen, Eberhard-Karls-Universität

<sup>2</sup>Klinik und Poliklinik für Mund-, Kiefer- und Gesichtschirurgie, Universitätsklinikum  
Tübingen

<sup>3</sup>Abt. für Radiologische Diagnostik, Radiologische Universitätsklinik, Universitätsklinikum  
Tübingen

#### **Purpose**

Skull bone tumours are highly complicated to resection especially those originating between external and internal skull table. The primary treatment is surgical resection. Traditional surgical approaches are associated with a significant bone deformity. We introduce and outline the clinical advantages of a navigation-assisted approach to lateral skull bone tumours using image-guided surgery.

#### **Material and Methods**

Our experience includes one patient with a bone tumour. Skull radiographic computed tomography revealed a bone lesion in the left temporo-parietal region. Computed tomographic scan indicated a destructive lesion involving external and internal skull bone layer. The patient was scheduled for image-guided surgery by use of a wireless passive infrared surgical navigation system (VectorVisionTM, BrainLAB). The preoperative computed tomography (CT) data was obtained before surgery using a newest generation Somatom Sensation 16 multi-slice-scanner (Siemens). After attaching the skull reference to the patients head, the patient-to-image registration was performed using laser surface scanning. The registration accuracy was expressed by a calculated value, the root mean

square (RMS) and while for system validation the intraoperative accuracy was visually checked by identification of anatomical landmarks.

Results: The procedure was successful and the tumour was minimally invasively removed, with no peri- and postoperative complications. The patient was ready for discharge 2 days after surgery. Postoperative imaging scans showed no recurrent tumour process.

## Conclusion

Traditional surgical approaches to skull bone tumours may result in significant cranial deformity and morbidity. Image-guided excision with surgical navigation techniques is a safe and effective minimally invasive surgical treatment option.